

# Analysis of the Level of Knowledge on the Tobacco in Adolescents from the Medical Sciences in Cuba

Carmen Cabelas Gonzalez<sup>1</sup>, Jesus Cuellar Alvarez<sup>2</sup>, Andrea Corsina Rodriguez<sup>3</sup>, Mirelys Olmo Quinones<sup>4</sup>

<sup>1</sup>Department of Medicine, Polyclinic “Jose Ramon Leon Acosta”, Cuba, <sup>2</sup>Department of Ceasing Tobacco, Polyclinic “Jose Ramon Leon Acosta”, Cuba, <sup>3</sup>Department of Nursing, Polyclinic “Jose Ramon Leon Acosta”, Cuba, <sup>4</sup>Department of Nursing, Polyclinic “Jose Ramon Leon Acosta”, Cuba

## ABSTRACT

**Background:** The analysis in adolescent’s smokers as prevention from the medical sciences is one of the lines of the work team in the consultation of ceasing tobacco. **Objective:** To analyze of the level of knowledge on the tobacco in adolescents from the medical sciences in Cuba. The investigation embraced one period from October 2018 to September 2019. **Method:** Was carried out a study decriptive, restrospective, for sampling intentional non probabilistic and with a universe composed by 18 old adolescents of the Polyclinic “Jose Ramon Leon Acosta”. It gathers it of the data was carried out through the empiric method as the clinical histories, interviews structured, the questionnaire and for the analysis of the data the statistical calculation was used. **Results:** The state of individual health of this group prevailed as for knowledge that will allow them to modify its lifestyle and in turn an aging very happened to the minimum of limitations. **Conclusion:** The difficulties are focused in the sistematic of the development psicosocial and to prevent risk factors in adolescents with addiction.

**Key words:** adolescence, addiction, model function, tobacco.

## INTRODUCTION

The adolescence is essentially a time of changes in which happens the process of the boy’s transformation in adult, has characteristic peculiar, and it is also a stage, of discovery of the own identity (psychological identity, sexual identity) as well as of the individual autonomy.

The adolescence like period of the human development has been object of the social scientists’ attention and of international instances who have tried to define its limits, as well as the characteristics that define this stage.<sup>1</sup>

The space of the adolescents has vital importance in the formative process of the personality, propitiating that the same ones can self-teach with a clearer vision of the harmful effects

that causes the tobacco and to prevent its consequences with the help of the professionals of the health but with a model function.

For model function understands each other those attitudes that impact of positive form or negative on the behaviors of the population’s health.<sup>2</sup>

The tobacco, is defined as a dysfunction caused by a substance able to produce dependence, the nicotine. It is considered a true drug diffused in the entire world. It reaches a world prevalence of 47 % in the masculine population of adults, in front of 12 % in the woman, and in the last years a more precocious beginning of the habit is observed, particularly in the beginning of the adolescence.<sup>3</sup>

### Address for correspondence:

Jesus Cuellar Alvarez, Department of Ceasing Tobacco, Polyclinic “Jose Ramon Leon Acosta”, Cuba.

DOI: 10.33309/2638-7719.050101

© 2022 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

The addiction of smoking could be immerse in the integral style of people's life, expert this as the group of individual decisions that affect the health and envelope which you could exercise certain control degree that also has an effect on the health and the behavior of those who cohabit. <sup>4</sup>

Almost 80 % of those more than a thousand million smokers that there is in the world lives in countries of low revenues or means, where it is bigger the mobility load and mortality associated to the tobacco. <sup>5</sup>

At the moment Cuba occupies the fifth place in Latin America and the Caribbean as for the prevalence of the tobacco, being Chile and Republic of the Dominican Republic the countries of more incidences in our continent. <sup>6</sup>

The phenomenon that exists around this real problem, conditioned the position of the following scientific problem in the present investigation: ¿How analyze the level of knowledge on the tobacco in adolescents from the medical sciences in Cuba?

General objective: to analyze of the level of knowledge on the tobacco in adolescents from the medical sciences in Cuba.

## METHODS

A study descriptive- retrospective, approach was carried out in adolescents patients attended from a community health area belonging to the "José Ramón León Acosta" policlinic of Santa Clara municipality, in the period from October 2018 to September 2019, with the objective of analyze of the level of knowledge on the tobacco in adolescents from the medical sciences in Cuba. The sample was selected through the simple random sampling probabilistic technique and previously informed consent to participate in the study.

It was study object an universe constituted by 18 adolescents and in those that was identified about of the level of knowledge on the tobacco in adolescents from a before and an after with risks related with the tobacco, to those which previously were requested informed consent.

## THEORETICAL LEVEL

□ Synthetic analytical: It made possible the interpretation of each one of the studied texts, to conform the criterion assumed in the epigraphs and paragraphs, as well as to particularize in the data obtained in the surveys to integrate them and to establish the corresponding generalizations.

□ Inductive-deductive: It facilitated going from the particular to the general in each of the analyzes carried out in the theoretical study and in the processing of the obtained information.

□ Generalization: It allowed the establishment of the regularities that were revealed in the study carried out.

## EMPIRIC LEVEL

□ Open interview: Contributed to identify the level of knowledge in the adolescents patients.

□ Individual clinical histories: It made it possible to provide information on various personal aspects.

□ Questionnaire: Contributed to identify the information on various personal aspects.

The selection was based on the following approaches: Inclusion approaches:

-All the adolescents with risks of the area of health that possess favorable psychic conditions to respond the questions.

-That they resided in the area of chosen health.

Exclusion approaches:

-Adolescents that emigrate of their residence place during the study.

Exit approaches:

-Adolescents that abandon the investigation voluntarily.

It was used the following variables starting from the obtained data: age and sex in the tobacco, and the level of knowledge on the tobacco in adolescents from a before and an after. Authorization was requested the adolescents, belonging to the educational policlinic "José Ramón León Acosta" for the realization of the study. To these they were explained the importance of the investigation. The data were used by the specialists of the health and with investigative ends, fulfilling the principle of the confidentiality of the data.

### Collection of the information

To begin the development of the investigation was carried out a bibliographical revision of the topic making a meticulous analysis of the most excellent aspects in the Cuban means as at international level. It was used as technical, the documental revision that included individual clinical histories and it was applied a questionnaire with the objective of obtaining information with respect to the level of knowledge on the tobacco in adolescents from a before and an after.

### Statistical prosecution

The information was stored in a file of data in SPSS version 21.0 and it is presented in statistical charts; for the description it was calculated the arithmetic stocking, standard deviation, absolute frequencies and percent's. In the analysis it was used non parametric tests as Squared Chi for adjustment kindness and independence of factors. One worked with significance levels to 5 %.

## RESULTS


The Chart 1 show ages that are located among the 12 years and more, 14 adolescents patients is of the male sex (77,7 %) and 4 of the female sex (22,2 %). In the analysis of the total of adolescents patients the sex male prevailed with 14 cases (77,7 %), the female one represents 4 woman (22,2 %), standing out the group of 12-15 years, for the two groups (77,7 %).

**Table 1: Analysis for sex and age**


Ages	Female		Male		Total	
	#	%	#	%	#	%
12-15	2	11,1	11	61,1	14	77,7
16-19	2	11,1	3	16,6	4	22,2
Total	4	22,2	14	77,7	18	100

Source: clinical history of health

Before the intervention, the basic knowledge on tobacco addiction are not high (56 %), after the application there was a change for well in the knowledge, until of a (83 %), what is translated in possible changes of lifestyles. The intervention improves the knowledge it has more than tobacco addiction after the intervention with a lifestyle good of more than 95 % (Fig. 2).



**Figure 1.** Level of knowledge on the tobacco in adolescents from a before



Source: clinical history of health

**Figure 2.** Level of knowledge on the tobacco in adolescents from an after

## DISCUSSION

The results of this investigation belonged together since with the world and national tendency every time it increases more the addiction in early ages. It coincides with a study carried out according to the authors<sup>7</sup> and with other investigations starting from authors like<sup>8</sup> on the tobacco in adolescents in a community in Spain, where it was reported that the half age of beginning in this habit, is located in the 13 years of age. A study in this respect having offered by another investigator as<sup>9</sup> they outline that the early beginning of the habit of smoking brings future problems of health and it is the entrance door for the consumption of alcohol or other drugs.

Due to the early age in that this addiction appears, was carried out this investigation where was appreciated that the committed ages in the study are from 12 to 15 years being predominant the masculine sex, coinciding with <sup>10-11</sup> that argue that the biggest prevalence in the tobacco is among the adolescents of the masculine sex, standing out the enormous risk that have of smoking the adolescents whose family makes it and the fateful consequences that brings the habit of smoking for its future life.

In Cuba the differences among the indexes of masculine and feminine adolescents' consumption are not marked, according to that referred in the National Program of Control of the Tobacco of the Ministry of Public Health outlined for<sup>12</sup> however doesn't behave of equal it forms study presently, where the masculine sex prevails on the feminine one.

In spite of the present behavior patterns in these adolescents, and the permissive of the habit for some parents and tutors, the way of obtaining of the cigarettes is through friends or contemporary, followed by traveling salespersons. This coincides with that found by other authors like<sup>14-15</sup>, although it is important to point out that in the study, a group of them obtains it in the schools and the practice of its habit is recurrent, that which is possible for the lack of professors' demand that facilitate the cigarette to the student in many of the cases. This belongs together with other studies like the one carried out in Argentina.<sup>16-17</sup>

On the other hand, other studies for <sup>18</sup>. they reflect the place of the smoking adolescents' more frequent consumption (50,9 %) between friends' house and parties, and in 40,4 % in their houses.

According to authors like <sup>19-20</sup> they outline that it is known the influence that has the example of the parents to adopt attitudes, by what becomes precise to surround the adolescents of positive examples that redound in the prevention of the tobacco, in order to reinforce the attitudes that go against the adoption of the habit of smoking, and where the family plays

a fundamental list, behaving as another factor of risk for the consumption of the tobacco in these precocious ages of the life.

In connection with the accessibility to the consumption of cigarettes stood out that in spite of the prohibition of sale of cigarettes to smaller than age in Cuba, 61 % of the interviewed active smokers acquire the cigarettes through the state stores or for traveling salespersons, without discarding that the road easier of obtaining it is by means.

## CONCLUSION

The male sex prevailed in the study, where the stadium average of beginning to present addiction to the tobacco is in adolescents. The magnitude of the population's aging doesn't have precedents; it is a process without limits in the humanity's history, the number of grown-ups increases exponentially in complex and uncertain socioeconomic joints. The development of professional competitions that offer attention to the adolescents to guarantee the quality and the excellence in the attention of health, will allow potenciar an appropriate lifestyle and to prevent that becomes a crisis factor for the sanitary structure. The obtained results are notably positive on the cost of their implementation in the current socioeconomic assisting to the focus psychologic and social in health, about the improvement of the lifestyle of this group.

### Conflict of interest

The author declares no conflict of interest.

## REFERENCES

- Gilbert Martínez C. Adolescencia y salud. Una mirada contemporánea. Ciudad México: Dirección Técnica de Servicios de Salud. Departamento de Medicina Preventiva; 2014: 15.
- Hurtado, M. Espacio sin humo Editorial José Martí, La Habana; 2016:
- Álvarez Valdés N, Gálvez Cabrera E, Díaz Garrido D. Hábito de fumar en la adolescencia al nivel comunitario. *Rev Cubana Med Gen Integr*; [ 2012; mayo; 17]; 23(3): Disponible en: [http://scielo.sld.cu/scielo.php?script=sci\\_arttext&pid=S0864-21252007000300006&lng=es](http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0864-21252007000300006&lng=es).
- Martínez E, Saldarriaga L. Hábito de fumar y estilo de vida en una población urbana. *Rev. Fac. Nac. Salud Pública* [internet]. 2011[citado mayo 2012]; 29(2): Disponible en: [http://www.scielo.org.co/scielo.php?pid=S0120-386X2011000200006&script=sci\\_arttext](http://www.scielo.org.co/scielo.php?pid=S0120-386X2011000200006&script=sci_arttext)
- Informe OMS sobre la epidemia mundial de tabaquismo, 2013 Disponible en: <http://www.who.int/mediacentre/factsheets/fs339/es/>
- Tabaquismo mata a más de 13 mil cubanos anualmente - Periódico La Demajagua, Diario digital de la provincia de Granma, Cuba Agencia Cubana de Noticias (ACN) | 31 mayo, 2016
- Ariza C, Nebot M, Villalbí JR, Díez E, Tomás Z, Valmayor S. Tendencias en el consumo de tabaco, alcohol y cannabis de los escolares de Barcelona. *Gac Sanit*. 2012; 17(3):190.
- Córdoba, R. Villalbí, R. Salvador, T y López, V. (2012). El proceso para la adopción de una legislación eficaz en la prevención del tabaquismo en España. *Rev Esp Salud Pública*: 80(4):631-45.
- Larmusch SY. Adolescencia y drogadicción. Un enfoque de la prevención escolar en Argentina. [Consultado 28 de julio de 2014]. Disponible en: <http://www.monografias.com/trabajos/adoldrog/adoldrog.shtml#top>
- El tabaco y los niños. American academy of child and adolescent psychiatry. [Consultado 28 de julio de 2014]. Disponible en: [http://www.aacap.org/cs/root/facts\\_for\\_families/informacion\\_para\\_la\\_familia/el\\_tabaco\\_y\\_los\\_ninos\\_no\\_68](http://www.aacap.org/cs/root/facts_for_families/informacion_para_la_familia/el_tabaco_y_los_ninos_no_68)
- Paterno CA. Factores de riesgo coronario en la adolescencia. Estudio FRICELA. [Consultado 31 de julio de 2014]. Disponible en: [http://www.revespcardiol.org/cgi-bin/wdbcgi.exe/cardio/mrevista\\_cardio.go?pid=13047009](http://www.revespcardiol.org/cgi-bin/wdbcgi.exe/cardio/mrevista_cardio.go?pid=13047009)
- Righetti J. Factores de riesgo en la niñez y adolescencia (Fundamentos de las recomendaciones FAC '99 en prevención cardiovascular). [Consultado 28 de julio de 2014]. Disponible en: <http://pcvc.sminter.com.ar/cvirtual/cvirtresp/cientesp/epesp/epc0024c/crighett/crighett.htm>
- Satcher D. Las mujeres y el fumar. Dirección General de Salud de los Estados Unidos. [Consultado 28 de julio de 2014]. Disponible en: [http://www.cancer.org/docroot/ESP/content/ESP\\_2\\_1x\\_Las\\_mujeres\\_y\\_el\\_fumar.asp?sitearea=ESP&viewmode=print&](http://www.cancer.org/docroot/ESP/content/ESP_2_1x_Las_mujeres_y_el_fumar.asp?sitearea=ESP&viewmode=print&)
- Achiong Estupiñán F. Prevalencia y riesgo atribuible al tabaquismo. *Rev Cubana Hig Epidemiol*. 2006;44(1):15.
- Montero Ramírez G. Hábito de fumar en adolescentes. *Educare* 21. Dic 2005 [Consultado: 2 de abril de 2015]; 8(3):15-8. Disponible en: <http://www.enfermeria21.com/educare/>
- Da Silva Pinto D, Aparecida Ribeiro S. Variables related to smoking initiation among students in public and private high schools in the city of Belém, Brazil. *J Bras Pneumol*. 2012;33(5):558-64.
- Bolzán A, Peleteiro R. Tabaquismo durante la adolescencia temprana. Estudio en escolares Argentinos. *J Pediatr*. 2013; 79(5):461-6.
- Verra F, Zabert G, Ferrante D, Morello P, Virgolini M. Consumo de tabaco en estudiantes de educación secundaria de Argentina. *Rev Panam Salud Publica*; [cited 2012 July 17]; 25(3): 227-233. Available from: [http://www.scielosp.org/scielo.php?script=sci\\_arttext&pid=S1020-49892009000300006&lng=en](http://www.scielosp.org/scielo.php?script=sci_arttext&pid=S1020-49892009000300006&lng=en)
- Ugarte Díaz RM. La familia como factores de riesgo, protección y resiliencia en la prevención del abuso de drogas en adolescentes. Capítulo 5. [Consultado 28 de julio de 2014].

[http://www.venumperu.com/friesgo\\_cap5\\_p130\\_169.pdf](http://www.venumperu.com/friesgo_cap5_p130_169.pdf)

20. González Henríquez I, Berger Vila K. Consumo de tabaco en adolescentes: factores de riesgo y factores protectores. [Consultado 28 de julio de 2014]. Disponible en: [http://www.scielo.cl/scielo.php?pid=S0717-95532002000200004&script=sci\\_arttext](http://www.scielo.cl/scielo.php?pid=S0717-95532002000200004&script=sci_arttext). ISSN 0717-9553.

**How to cite this article:** Gonzalez C C, Alvarez J C, Rodriguez A C, Quinones M O. Analysis of the Level of Knowledge on the Tobacco in Adolescents from the Medical Sciences in Cuba. *J Community Prev Med* 2022;5(1):01-05.  
DOI: 10.33309/2638-7719.050101