

A Girl from Canada with Severe Cerebral Palsy Associated with Hydrocephalus and Mutation of Kinase D-Interacting Substrate of 220-KDa (KIDINS220) Gene: A New Syndrome with Unique Brain Imaging Findings and a Therapeutic Challenge

Aamir Jalal Al-Mosawi

Advisor and expert trainer, The National Training and Development Center, Iraqi Ministry of Health, Baghdad, Iraq

ABSTRACT

Background: Cerebral palsy is a heterogeneous neurologic condition caused by abnormalities of brain development or damage affecting mostly regions controlling movements, balance, and posture. The brain abnormalities and damage causing cerebral palsy are characteristically non-progressive, but lead to permanent abnormalities of movements, posture, and limitation of mobility. Kinase D-interacting substrate of 220 kDa (KIDINS220), a trans-membrane protein that is essential for growth mediating pathways in the nervous system. Heterozygous KIDINS220 truncating variants affecting protein's C-terminus have been reported only in few unrelated children who had spastic paraplegia, mental retardation, nystagmus, and obesity. A homozygous, more N-terminal truncating variant in KIDINS220 gene has been reported to be associated with enlarged brain ventricles and limb contractures in three fetuses from a consanguineous family. **Patients and methods:** The family of a girl who developed severe cerebral palsy caused by hydrocephalus, and had mutation of kinase D-interacting substrate of 220-kDa (Kidins220), consulted us about the possible therapies her condition. Unique brain imaging abnormalities are described, and the relevant literatures were reviewed with aim of suggesting possible evidence-based therapies. **Results:** A 33-month old girl from Canada had cerebral palsy dominated by spasticity and caused hydrocephalus. The girl had slight nystagmus, and was also thought to experience sometimes dystonia. During the first 18 months of life, her head was enlarging, and Brain MRI showed dilated lateral ventricles, hypoplastic basal ganglia and inferior vermis, agenesis of septum pelucidum, thin corpus collosum. Before surgery (endoscopic third ventriculostomy) at 19 months of age, her head circumference was above the 99th percentile. Her head stabilized after the surgery. Literature review suggested the possible usefulness of the use multi-factorial therapies including cerebrolysin, citicoline, piracetam, and pyritinol, and based on our extensive clinical experience we suggested an initial one-month therapeutic course. **Conclusion:** In this paper, a new genetic syndrome is described. The syndrome is associated with mutation of KIDINS220 gene, cerebral palsy, hydrocephalus, nystagmus, and unique brain imaging findings including hypoplastic basal ganglia and inferior vermis, agenesis of septum pelucidum, and thin corpus collosum.

Key words: New genetic syndrome, mutation of KIDINS220 gene, cerebral palsy, hydrocephalus, nystagmus.

Address for correspondence:

Aamir Jalal Al-Mosawi, Advisor and expert trainer, The National Training and Development Center, Iraqi Ministry of Health. Baghdad, Iraq.

DOI: 10.33309/2639-913X.040105

© 2021 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

INTRODUCTION

Cerebral palsy is a heterogeneous neurologic condition caused by abnormalities of brain development or damage affecting mostly regions controlling movements, balance, and posture. The brain abnormalities and damage causing cerebral palsy are characteristically non-progressive, but lead to permanent abnormalities of movements, posture, and limitation of mobility. [1, 2, 3, 4].

We reported the occurrence of various brain imaging abnormalities in children with cerebral palsy including brain atrophy, periventricular leukomalacia, arachnoid cyst, and cerebellar atrophy/cerebellar hypoplasia [5-15].

Kinase D-interacting substrate of 220 kDa (KIDINS220), a trans-membrane protein that is essential for growth mediating pathways in the nervous system. Heterozygous KIDINS220 truncating variants affecting protein's C-terminus have been reported only in few unrelated children who had spastic paraplegia, mental retardation, nystagmus, and obesity.

A homozygous, more N-terminal truncating variant in KIDINS220 gene has been reported to be associated with enlarged brain ventricles and limb contractures in three fetuses from a consanguineous family [16].

PATIENTS AND METHODS

The family of a girl who developed severe cerebral palsy caused by hydrocephalus, and had mutation of kinase D-interacting substrate of 220-kDa (Kidins220), consulted us about the possible therapies her condition. Unique brain imaging abnormalities are described, and the relevant literatures were reviewed with aim of suggesting possible evidence-based therapies.

RESULTS

The 33-month old girl from Canada was studied during October, 2021 (Figure-1). She had cerebral palsy dominated by spasticity and caused hydrocephalus. She had significant delay in gross motor, fine motor, speaking, and was unable to sit unassisted and had poor head control. She was unable to neither roll, crawl, sit, nor stand. Spasticity had already associated with flexion deformity at the ankle which was correctable. However, the child was able to recognize the face of the mother, and was smiling, and responding to her name. She was starting to make some sounds with words, na, ma. She also had slight nystagmus. The girl was also thought to experience sometimes dystonia.

During the first 18 months of life, her head was enlarging, and Brain MRI showed dilated lateral ventricles, hypoplastic basal ganglia and inferior vermis, agenesis of septum pelucidum, thin corpus callosum



Figure 1. 33-month old girl from Canada who had cerebral palsy dominated by spasticity

Before surgery (endoscopic third ventriculostomy) at 19 months of age, her head circumference was above the 99th percentile. Her head stabilized after the surgery.

She was receiving baclofen 10 mg at bed time, gabapentin, and omeprazole for gastro-esophageal reflux.

Literature review suggested the possible usefulness of the use multi-factorial therapies including cerebrolysin, citicoline, piracetam, and pyritinol, and based on our extensive clinical experience we suggested an initial one-month therapeutic course [1-15].

The Initial Therapeutic Course Primarily Aimed at

1-Improving posture and spontaneous movement by providing adequate muscle relaxation which hopefully results in easier correction of the flexion deformity at the ankle.

2-Repairing brain which can result in improved brain function that can be manifested early by improvement in head control, and sometimes may result an early cognitive improvement which can be associated with improved speech and a better understanding of simple speech. Improvement in nystagmus may occur early.

Our suggested evidence-based treatment included withdrawal of gabapentin, and the gradual increase of baclofen to 20 mg daily in 4 divided doses during the first month of treatment.

Initially, Baclofen 5 mg is given three times a day, and increased to 4 times a day on day-8. Diazepam 2 mg daily at bed time is added at day 14, and continued.

In addition, Cerebrolysin 2 ml given by intramuscular injection daily every other day during the morning hours (15 doses over one month). Oral Citicoline syrup 2 ml (200mg) daily in the morning was also prescribed.

DISCUSSION

Neuroimaging studies showing enlarged ventricles suggest the diagnosis of hydrocephalus in the presence of head enlargement. However, as early as 1996, the work of Campistol and colleagues suggested that in the absence of macrocephaly, brain images showing slight ventricular dilation, normotensive hydrocephalus, hydrocephalus with irregular limits, subcortical atrophy or periventricular heterotopia may indicate residual periventricular leukomalacia [17].

From Spain reviewed 250 patients with cerebral palsy, and reported their brain imaging findings which included atrophy in 38.8%, hydrocephalus in 29.4%, ischemia in 14.9%, hemorrhage in 11.6%, and no abnormalities in 23.8% [18].

From the United Kingdom reviewed patients with cerebral palsy born in 1966-1991 in the counties of Merseyside and Cheshire, UK, and reported an important association between cerebral palsy and hydrocephalus [19].

Reviewed 1104 patients (663 males, 441 females) with cerebral palsy were born during the period from 1991 to 1999 in three areas (Isère Region, French Alps; Funen County, Denmark; Northern Region, England). They found that the most common cerebral anomalies were primary microcephaly which was observed in 26.5%, and congenital hydrocephalus which was observed in 17.3% [20].

Reported brain MRI and CT-scan findings in 129 children (86 males, 43 females; aged 2-18 years) who had various types of cerebral palsy. 100 patients (77.52%) had CT-scan and 29 patients (22.48%) had MRI. 19 patients (14.72%) had normal MRI and CT-scan. Ventriculomegaly which indicates hydrocephalus in the presence of head enlargement was the most common findings and occurred in 38 patients, followed by focal infarct which was observed in 28 patients, periventricular leukomalacia was observed in 17 patients, cortical and cortical/subcortical atrophy was observed in 13 patients, basal ganglia lesions was observed in 12 patients, and cortical migration and organization problems were observed in 12 patients [21].

Kinase D-interacting substrate of 220 kDa (KIDINS220), a trans-membrane protein that is essential for growth mediating pathways in the nervous system. Heterozygous KIDINS220 truncating variants affecting protein's C-terminus have been reported only in few unrelated children who had spastic paraplegia, mental retardation, nystagmus, and obesity.

A homozygous, more N-terminal truncating variant in KIDINS220 gene has been reported to be associated with enlarged brain ventricles and limb contractures in three fetuses from a consanguineous family [16].

CONCLUSION

In this paper, a new genetic syndrome is described. The syndrome is associated with mutation of KIDINS220 gene, cerebral palsy, hydrocephalus, nystagmus, and unique brain imaging findings including hypoplastic basal ganglia and inferior vermis, agenesis of septum pelucidum, and thin corpus callosum.

ACKNOWLEDGEMENT

The author would like to express his gratitude for the family of the patients who willingly accepted publishing the photo of their child.

REFERENCES

1. Al Mosawi AJ. Dramatic effect of nandrolone decanoate on motor development in cerebral palsy. Arch Dis Child 2008; 93(11) :(Suppl II):A338-A339.
2. Al Mosawi AJ. A novel therapeutic approach for the treatment of cerebral palsy (Ed). LAP Lambert Academic Publishing, Germany, 2017.
3. Al Mosawi AJ. A novel therapeutic approach for the treatment of brain atrophy (Ed). LAP Lambert Academic Publishing, Germany, 2017.
4. Al Mosawi AJ. New therapies for the treatment of spastic cerebral palsy (Ed). LAP Lambert Academic Publishing, Germany, 2019.
5. Al-Mosawi AJ. New therapies for the treatment of spastic cerebral palsy. Med J Clin Trials Case Stud (ISSN: 2578-4838) 2019; 3(2): 000209. USA. DOI: 10.23880/mjccs-16000209.
6. Al-Mosawi AJ. The pattern of cerebral palsy in Iraqi children. Med Life Clin. 2019; 1(1):1001.
7. Al-Mosawi AJ. The pattern of cerebral palsy in Iraqi children. 1st ed., Saarbrücken; LAP Lambert Academic Publishing: 2019 (ISBN: 978-620-0-09427-8).
8. Al-Mosawi AJ. New Therapies for the treatment of ataxic cerebral palsy caused by kernicterus. EC Clinical and Medical Case Reports March 18, 2020; 3(4): 26-31.Doi: 10. 5281/zenodo.3892208.

9. Al-Mosawi AJ. The experience with the use of nandrolone decanoate and pyritinol in children with cerebral palsy. *Open Access Journal of Biogeneric Science and Research* (ISSN: 2692-1081) July 6, 2020; 2(3):1-3 Doi: 10.46718/JBGSR.2020.01.000 051.
10. Al-Mosawi AJ. Cerebral palsy: A unique illustrated experience. *Medico Research Chronicles* (ISSN: 2394-3971) 2020 September 8; 7 (4):2017-239. Doi: 10.26838/MEDRECH.2020.7.4.441.
11. Al-Mosawi AJ. Congenital Externally Communicating Porencephaly Presenting as Hemiplegic Cerebral Palsy: Imaging Study of a Rare Condition. *SunKrist Journal of Neonatology and Pediatrics* January 18, 2021; 3 (1):1-4. Doi: 10.5281/zenodo.4515510.
12. Al-Mosawi AJ. The early treatment of a boy from Virginia with ataxic cerebral palsy. *Journal of Pediatrics and Child Health* 2021 May 22; 2 (4): 1-5. Doi: 10.5281/zenodo.4777413 .
13. Al-Mosawi AJ. Cerebral palsy: An illustrated ground-breaking experience. *Scholar's press: 2020* (ISBN: 978-613-8-93772).
14. Al-Mosawi AJ. Mózgowe porażenie dziecięce: Ilustrowane przełomowe doświadczenie: Medycyna oparta na dowodach (Polish Edition). *Wydawnictwo Nasza Wiedza* September 16, 2020 (ISBN-10: 6202762136, ISBN-13: 978-620 276 2 137).
15. Al-Mosawi AJ. Cerebrale parese: Een geïllustreerde baanbrekende ervaring: Bewijsgerichte geneeskunde (Dutch Edition) .*Uitgeverij Onze Kennis : September 16, 2020* (ISBN-10 : 6202762128, ISBN-13 : 978-6202762120).
16. El-Dessouky SH, Issa MY, Aboulghar MM, Gaafar HM, Elarab AE, Ateya MI, Omar HH, Beetz C, Zaki MS. Prenatal delineation of a distinct lethal fetal syndrome caused by a homozygous truncating KIDINS220 variant. *Am J Med Genet A* 2020 Dec; 182 (12): 2867-2876. Doi: 10.1002/ajmg.a.61858. PMID: 32909676.
17. Campistol Plana J, Escofet Soteras C, Póo Argüelles P. Leucomalacia periventricular: diagnóstico retrospectivo en niños con diplejía espástica [Periventricular leukomalacia: its retrospective diagnosis in children with spastic diplegia]. *An Esp Pediatr* 1996 Jun; 44(6):553-6. PMID: 8849097 [Article in Spanish].
18. Bringas-Grande A, Fernández-Luque A, García-Alfaro C, Barrera-Chacón M, Toledo-González M, Domínguez-Roldá JM. Parálisis cerebral infantil: estudio de 250 casos [Cerebral palsy in childhood: 250 cases report]. *Rev Neurol* 2002 Nov 1-15; 35(9):812-7. PMID: 12436377 [Article in Spanish].
19. Pharoah PO. Prevalence and pathogenesis of congenital anomalies in cerebral palsy. *Arch Dis Child Fetal Neonatal Ed* 2007 Nov; 92(6):F489-93. Doi: 10.1136/adc.2006.107 375. PMID: 17428819.
20. Rankin J, Cans C, Garne E, Colver A, Dolk H, Uldall P, Amar E, Krageloh-Mann I. Congenital anomalies in children with cerebral palsy: a population-based record linkage study. *Dev Med Child Neurol* 2010 Apr; 52(4):345-51. Doi: 10.1111/j.1469-8749.2009.03415.x. PMID: 19737295.
21. Grigore I, Diaconu G. Corelații clinico-radiologice în paralizia cerebrală [Clinical and radiologic correlations in cerebral palsy]. *Rev Med Chir Soc Med Nat Iasi* 2010 Jul Sep; 114(3):748-52. PMID: 21243802 [Article in Romanian].

How to cite this article: Al-Mosawi AJ. A Girl from Canada with Severe Cerebral Palsy Associated with Hydrocephalus and Mutation of Kinase D-Interacting Substrate of 220-KDa (KIDINS220) Gene: A New Syndrome with Unique Brain Imaging Findings and a Therapeutic Challenge. *J Clin Res Radiol* 2021;4(1):22-25. DOI: 10.33309/2639-913X.040105